Animation of Flocks Flying in Line Formations

Marina Klotsman & Ayellet Tal Department of Electrical Engineering, Technion

Corresponding author: Ayellet Tal ayellet@ee.technion.ac.il, phone: +972-4829-4651

Abstract

The coordinated flight of bird flocks is a pleasant and attractive sight. We provide a biologicallymotivated technique for modeling and animating bird flocks, which produces plausible and realisticlooking flock animations. While most previous approaches have focused on animating cluster formations, this paper introduces a technique for animating flocks that fly in certain patterns, so-called line formations. We distinguish between the behavior of such flocks during initiation and their behavior during steady flight. Our simulation of the initiation stage is rule-based and incorporates an artificial bird model. Our simulation of the steady-flight stage combines a data-driven approach and an energy-savings model.

Keywords: artificial flocks, artificial birds, animation.

I. INTRODUCTION

Researchers in biology and computer graphics have long been intrigued by the challenge of realistically modeling and animating groups of animals. One such example is the synchronized motion of flocks of birds, which is a delightful and fascinating sight. Flying flocks can be classified by their formation – *cluster formations* or *line formations* [10]. In cluster formations, typical of small birds as well as fish and herds, the animals are organized in irregular shapes. Conversely, in line formations the flocks are organized in a characteristic pattern, such as a line, 'V', or 'U'. These formations are typically two-dimensional and exhibit a high degree of regularity in spacing and alignment. They are common in flocks of large birds, such as waterfowls, cranes, and pelicans.

In his ground-breaking paper, Reynolds proposed animating groups by simulating the behavior of each bird independently [24]. The results are pretty animations of cluster flocks. Others have also addressed the animation of cluster formations [12], [4]. This paper introduces a technique for realistically animating flocks of birds flying in line formations (Figure 1).

In nature, these flocks undergo two stages: flock *initiation* (also termed *formation*) and *steady flight*. Initiation occurs during takeoff, with many rapid changes in the locations of the birds within the flock. During steady flight the flock flies over large distances in more stable shapes. Previous approaches have not distinguished between these stages and applied the same rules to the whole animation. While this strategy suffices for clusters, it does not suit the long, steady flights of flocks flying in line formations. We propose modeling these stages differently.

Why flocks fly long distances (up to 20,000 miles) in specific patterns is still a puzzle [2]. While it is often believed that the central rationale is to save energy [32], no model exists that can precisely predict the birds' positions. This lack of knowledge directed us into choosing a data-driven approach, utilizing real examples of flying flocks, for animating the steady-flight



Fig. 1. Animation of a flock of barnacle geese (snapshots)

3

stage. Since existing examples are typically very short, while the flight is extremely long, we augment these examples with calculations of potential energy-savings. This combination is shown to produce not only feasible, but also eye-pleasing animations of flying bird flocks.

Inspired by previous work, our approach for animating the initiation stage is rule-based, where the behavior of each individual bird governs the motion of the whole flock. We introduce a new set of rules that better suits line formations. These rules are based on two novel drives, targetreaching, and separation, which help us more accurately animate the flock in initiation.

Our contribution is hence threefold. First, we propose to distinguish between the initiation and the steady flight (Sec. III). Second, we show how steady flight patterns can be produced by combining a data-driven approach with an energy-savings model (Sec. V). Finally, we suggest a novel set of rules for the initiation stage of line-formation flocks (Sec. VI).

In http://www.ee.technion.ac.il/ ayellet/Movies/10-KlotsmanTal.mpg, a video that demonstrates our results can be viewed.

II. RELATED WORK

Most previous work on flock animation has focused on flocks flying in clusters. We briefly introduce this work. Then we present the prevailing biological hypotheses that attempt to explain why certain flocks fly in line-formation patterns and a couple of papers that animate such flocks. **Cluster formations:** Animation of group behavior – of humans and animals alike – has attracted a lot of attention in computer graphics [17], [18], [21], [24], [25], [27], [29], [30], [31]. We focus on bird flocks.

Heppner defines a *flock* as a group of flying birds characterized by coordination in one or more of the following flight parameters: turning, spacing, timing of takeoff and landing, and individual flight speed and direction [10]. In his seminal work, Reynolds proposes three rules for animating flocks: collision avoidance, velocity matching, and flock centering [24]. The aggregate flock motion is a result of the actions of all the birds. In [12], stochastic differential equations are developed on the basis of four drives: homing. interaction, attempting to preserve a specific flight velocity, and randomness. In [23], it is proposed to model flocks by non-linear dynamics. In [1] a technique is described for generating constrained group animations. In [4], fuzzy logic is introduced to flock modeling. This allows the model to use uncertain knowledge

and linguistic descriptions. The results are estimated and shown to improve the results of [24] in some parameters.

These techniques exhibit pretty animations for cluster-shaped flocks. However, line-formation flocks are difficult to generate using these methods, since the rules employed tend to "cluster" the birds together. In [4], specific, but unrealistic, initial conditions may produce line formations, but these are likely to break during the flight.

Line formations: There are two main hypotheses explaining the tendency of birds to adopt a line formation: enhanced communication and aerodynamic energy savings [2]. According to the first hypothesis, these formations are the result of the visual requirements of line-formation flight. According to the second, a bird in a line formation can save energy by taking advantage of the wingtip vortex produced by the neighboring birds [11].

Without taking into account wing flapping, [19] predict the optimal distances between the birds, which yield maximal energy savings. A specific "U" shape is shown to save energy equally for all the birds in the flock. In [13], [14], [28], [26], more elaborate and accurate power-reduction calculations are presented. [2] suggest that small flocks containing related birds use "V" formations, where the leading birds save less energy.

Field studies, however, show a wide variation in spacing and in deviations from the above optimal positions. Some interpret these deviations as a failure of the birds to reach an optimal energy distribution [3], [9], while others view it as support for the alternative hypothesis [5], [8]. In [32], empirical evidence supports the energy-savings hypothesis: the heart rates of white pelicans are measured and significant energy savings shown for flocks flying in line formations. We adopt the energy-savings model, which is currently the dominant hypothesis.

In [7] it is proposed to add an additional rule to [24] in order to animate line-formation flocks. This rule keeps some portion of the bird's field of view unobstructed. This causes some of the birds to arrange into fragments of linear formations. Yet, such arrangements are local and the convergence of the whole flock to a line formation is not guaranteed. In [22], three rules are introduced for animating the initiation stage: coalescing (seeking proximity to the nearest bird), gap-seeking (seeking the nearest position with an unobstructed view), and stationing (avoiding downwash regions). It is demonstrated that the flock converges to a line formation. However, the realism of the animation is not guaranteed, since the directions of the birds are not taken into account (e.g., a bird is assumed to fly forward, even when its position changes horizontally).

We propose to solve these problems by providing a new set of rules for simulating the initiation and a novel data-driven approach, based on an energy-savings model, for animating steady flight, which has not been addressed before.

III. GENERAL APPROACH

In nature, during initiation and takeoff the positions and the velocities of the birds change rapidly. In contrast, smaller changes occur during the almost two-dimensional steady flight [10]. Our goal is to animate these different behaviors, so that the generated animations resemble flocks in nature.

Since no model exists that accurately predicts the behavior of a flock, we turn to a data-driven approach, i.e., learning from examples. Given an example movie, the challenges are to determine the major variables that characterize a given flock, extract the parameters that are relevant for computing them, compute these variables, and utilize them in the creation of plausible flock shapes.

While data-driven methods have been used successfully in crowd simulation [17], [18], [20], they cannot be used in a straightforward manner in our case. This is because, unlike human pedestrians who have their own individual trajectories, a bird flock has a common global goal—migrating together. Thus, while pedestrian behavior can be learned from various scenes in different movies, the trajectories of birds must maintain the sequential ordering of the original movie. Using the shapes in a single movie is unsatisfactory as well, since movies are typically very short and contain a limited number of possible flock shapes.

Hence, our solution combines the data-driven approach with an energy-savings model. The key idea is to learn from examples—but to learn the energy-savings (power reduction) values—not the trajectories, since these variable are generally believed to be the primary reason for flying in a line-formation pattern. A bird flying alone at steady speed demands more flight power than a bird flying in a flock, where drag reduction helps the bird save energy. *Power reduction* is defined as the ratio between this difference in flight power, that of a single bird minus that of a bird in a flock, and the flight power of a single bird.

We aim at generating shapes that preserve similar power-reduction values as those obtained in the example shapes, thus guaranteeing plausible shapes. Another benefit of this scheme is that



Fig. 2. Initiation of a "V" shape. The initiation begins with birds distributed randomly (left) and ends when the flock converges to the target shape (right). The arrows represent the flight directions of the birds.

it does not constrain us to the existing example shapes, in terms of the number of birds in the flock, the relative location of the leader bird, and the variations in birds' positions.

Alternative simpler methods, such as interpolation, not only limit the number of possible shapes (e.g., the shapes in Figure 4 are not interpolations of the given shapes), but might also generate shapes that do not comply with feasible energy-savings values, and thus could not occur in nature.

Below we describe the three components of our method: parameter extraction, steady flight, and flock initiation. We elaborate in Sections IV–VI. **Parameter extraction:** Most existing movies of line-formation flocks show small portions of the flight in a formation (steady flight). We aim at extracting the parameters that can help us compute the power-reduction values of the birds, which are essential in creating plausible flock shapes. These parameters are the flock's optimal shapes and outliers.

Steady flight: In nature, the shape of a flock changes constantly, and an optimal shape, as predicted by known models of energy-savings, is seldom obtained. Therefore, our approach incorporates energy-savings values with a data-driven approach.

Initiation: The goal is to produce an animation that begins with a given (e.g., random) distribution of the birds and ends when the flock converges to the first shape in the steady-flight animation (Figure 2). As in previous *artificial bird* models, our animated bird lives in a *world* and *perceives* its state through its senses. The bird's *actions* are governed by its interpreted perception and attempt to satisfy its *drives*. We propose a new set of drives, suitable for flocks flying in line formations: target-reaching (the tendency of a bird to reach a specific distance from a chosen reference bird) and separation (the tendency of a bird to have a unique reference

(a) U shape (b) J shape (c) V shape (d) Line shape

Fig. 3. Examples of flock shapes

bird to follow). Figure 3 shows some examples of how these rules, when combined, model the convergence of the flock to some shapes.

IV. PARAMETER EXTRACTION

We are given a movie of a real \hat{N} -bird flock flying in a formation (steady flight). Our goal is to extract the parameters that would let us compute the power-reduction values that would be later used to produce the animation. The algorithm consists of two steps. First, the position parameters are extracted for each movie frame. Then, the optimal geometric shape of the flock and the outliers are computed for each frame.

Step 1 – Position parameter extraction: The birds are first sorted by their horizontal (x) positions in the first frame and indexed accordingly. Then, the following parameters are extracted for each frame f:

- 1) Ordering: The order of the birds according to their *x*-positions.
- 2) *x*-distance statistics: The mean μ_{xf} and the standard deviation σ_{xf} of the horizontal distances between neighboring birds.
- 3) Leader bird: The index of the leading (front-most) bird.

Step 2– Computing the optimal shape and the outliers: To compute the optimal geometric shape that matches the flock's shape in a given frame and its outliers, we use the RANSAC (Random Sample Consensus) algorithm [6]. RANSAC allows a model to be fit to a data set that contains outliers and these outliers to be detected. It achieves its goal by iteratively selecting a random subset of the original data. This subset, composed of hypothesized inliers, is then tested. This procedure is repeated, each time producing either a model that is rejected because too few points are classified as inliers, or a refined model together with a corresponding error measure.

In our case, the input to RANSAC is the positions and the possible models. These models are either a parabola for $\{U,J\}$ -shapes or a linear function for $\{V,line\}$ -shapes. The output is the



Fig. 4. Shapes produced by our algorithm, given an example movie of 16 birds. The left two images show two different shapes of a flock consisting of 16 birds, whereas the right images show a flock of 28 birds. Note that the shapes are similar, but the sides are reversed. These generated shapes differ from the input movie, yet have similar energy-savings values.

model's parameters and the model's outliers. At the end of this stage, we produce the following information:

- 1) The optimal geometric shape that best fits the current formation, as well as the outliers.
- 2) Position deviations: The inliers' mean of the y-positions μ_{yf} and the standard deviation σ_{yf} .
- 3) Special roles: Being either a leader or an outlier bird is considered a special role.

V. STEADY FLIGHT

In nature, the steady flight stage is very long and is characterized by similar velocities of the birds and small frequent changes in their relative positions. Our proposed solution combines an energy-savings model with a data-driven approach. As illustrated in Figure 4, it does not constrain us to the existing example shapes, in terms of the number of birds, the relative location of the leader bird, and the variations in birds' positions.

The input consists of the parameters extracted (Section IV), the physical characteristic of the bird in terms of the wing-span b, the required number of birds in the animation N, and the required number of frames in the animation F. The goal is to construct an animation of a steady flight of a flock.

The key idea is to learn the flock's energy-savings patterns and generate shapes that respect them, thus creating plausible shapes. An important and novel aspect of our scheme is showing how the positions of the birds can be computed by solving a system of non-linear equations that take the power reduction values into account.

The algorithm, which is outlined in Algorithm 1, proceeds in four steps. We first create some initial shapes whose geometry is similar to the shapes in the example movie, but whose number of birds is N. These shapes are used to extract the range of power reduction values for each

bird. These values allow us to generate a wider variety of plausible shapes that comply with the energy-savings pattern. Once these plausible shapes are produced, they are interpolated to generate the final animation.

Algorithm 1 Steady flight animation Input:

Parameters extracted in the previous sections (shapes, deviations, special roles);

b – wing-span;

N -#birds in the animation;

F – #frames in the animation.

Output: An animation (*F* frames) of an *N*-bird flock, which consistes of of $F \times N$ position vectors.

Algorithm:

- 1) Construct initial shapes of an *N*-bird flock, which maintain the general shapes and deviations of the movie.
- 2) Calculate the power-reduction values of each bird, for each shape constructed in Step 1.
- 3) Construct new key shapes that satisfy the power-reduction values calculated in Step 2.
- 4) Create an *F*-frame animation of a steady flight, by interpolating between the shapes generated in Steps 1 and 3.

Step 1 – Constructing initial shapes: Initially, the first frame is used to derive a correspondence between the animated birds and the movie birds. This is done by first generating a geometric curve that maintains the frame's shape parameters. The required number of birds are distributed uniformly on this geometric curve, but the leader's relative position may differ, so as to suit its relative position in the initiation. The correspondence between the birds is then performed as follows: After matching the leaders—the front-most bird of the movie and the front-most bird of the animation (irrespectively of their relative location)—the sides containing more birds are matched. This is important, since each side in a line formation may have different shape parameters. Next, the birds closest to the leader and farthest from it (on each side) are matched, followed by matching the outlier birds, then the inliers, and finally adding $N - \hat{N}$ birds if necessary (uniformly distributed). Then, for every movie frame an animation frame is generated, as follows. The geometric curve that maintains the same parameters computed in Step 2 of Section IV is used. We need to find the exact locations of the birds on this curve. We first generate new *x*-distances, by drawing a value from a Gaussian distribution, having the extracted mean μ_{xf} and standard deviation σ_{xf} . Then, the initial *y*-position of each animated bird is calculated according to curve's parameters. Next, given the correspondence between every animated bird to a movie bird, the *y*-positions of birds whose corresponding example birds are outliers are modified, by adding the appropriate deviations. This is important in order for a flock to look "real." Finally, we add deviations to all other birds, which are drawn from a Gaussian distribution with μ_{yf} and σ_{yf} .

Step 2 – Power-reduction calculation: For each given shape produced in Step 1, we compute the power reduction value of each bird in the flock. We use the model of [13], which assumes a two-dimensional flock, with equal velocities and non-flapping wings. (More accurate models do not result in an analytic formula [16], [28].)

A flying bird creates vortices resulting in regions of *upwash* and *downwash* behind it, as illustrated in Figure 5. Other birds flying in this upwash region save energy by a drag-force reduction. Flying in a downwash region increases the drag and therefore is not beneficial. Each bird is affected by the upwash field generated by all the other birds in the flock, where the influence of laterally-distant birds is small.



Fig. 5. The downwash and upwash regions behind by the bird. The gray area is the downwash region, whereas the white area behind the bird is the upwash region.

Let C be the drag ratio and g_{ij} be a function below, which depends on the distance between birds *i* and *j*. When the birds in the flock have similar body parameters, the power reduction for bird *i* is

$$e_i = C \sum_{j=1}^N g_{ij}.$$
 (1)

Since the drag ratio C is constant for all the birds under our assumptions, we ignore it from now on.

Let x_{ij} $(i \neq j)$ be the lateral distance between the center of bodies of birds *i* and *j* and y_{ij} be their vertical distance, normalized by the wingspan *b*. y_{ij} is positive if bird *i* is behind bird *j* and negative otherwise. Assuming that the birds do not fly in the downwash regions, g_{ij} is given by [13]:

$$\frac{\pi^2}{2}g_{ij} = \ln\frac{x_{ij}^2}{(x_{ij} - \frac{\pi}{4})(x_{ij} + \frac{\pi}{4})} + 2\ln\frac{x_{ij} - y_{ij} + \sqrt{x_{ij}^2 + y_{ij}^2}}{x_{ij} + y_{ij} + \sqrt{x_{ij}^2 + y_{ij}^2}} - \ln\frac{x_{ij} - \frac{\pi}{4} - y_{ij} + \sqrt{(x_{ij} - \frac{\pi}{4})^2 + y_{ij}^2}}{x_{ij} - \frac{\pi}{4} + y_{ij} + \sqrt{(x_{ij} - \frac{\pi}{4})^2 + y_{ij}^2}} - \ln\frac{x_{ij} - \frac{\pi}{4} - y_{ij} + \sqrt{(x_{ij} - \frac{\pi}{4})^2 + y_{ij}^2}}{x_{ij} - \frac{\pi}{4} + y_{ij} + \sqrt{(x_{ij} - \frac{\pi}{4})^2 + y_{ij}^2}} + \frac{1}{y_{ij}} \left[2\sqrt{x_{ij}^2 + y_{ij}^2} - \sqrt{(x_{ij} + \frac{\pi}{4})^2 + y_{ij}^2} - \sqrt{(x_{ij} + \frac{\pi}{4})^2 + y_{ij}^2} \right].$$
 (2)

In this function the power reduction decreases non-linearly to zero as the x-distances increase and increases to a constant value as the y-distances increase, as illustrated in Figure 6.



Fig. 6. The geometric function g_{ij}

Step 3 – Constructing feasible key shapes: The goal is to construct a variety of feasible key shapes, which satisfy the movie's power-reduction values, but may differ from the original movie in shape. This is performed by first associating every animated bird with a range of "typical" power-reduction values (obtained at Step 2) and then generating new shapes that comply with this range.

In fact, we are interested in the relative power reduction values and not the absolute ones, since the relative values characterize the flock's shape and are influenced less by the exact lateral distances between the birds.

Let *E* be the total power reduction of the entire flock $E = \sum_{i=1}^{N} e_i$. Given e_i (Equation 1), the relative value of bird *i* is defined by:

$$\xi_{i} = \frac{e_{i}}{E} = \frac{e_{i}}{\sum_{n=1}^{N} e_{n}} = \frac{\sum_{j=1}^{N} g_{ij}}{\sum_{n=1}^{N} \sum_{j=1}^{N} g_{nj}}.$$
(3)

Let P_i be the maximal value of ξ_i over all the frames and p_i be the minimal value. A feasible power reduction range for bird *i* is defined by $[(1 - \varepsilon) \cdot p_i, (1 + \varepsilon) \cdot P_i]$.

To generate key shapes for the animation, we want to produce values that are not only within this range, but also guarantee smooth transitions between the key shapes. Thus, to generate a key shape k, a new relative power reduction value $\hat{\xi}_{i,k}$ is generated by randomly choosing a value in a range:

$$\hat{\xi}_{i,k} \in [(1-\varepsilon) \cdot p_i, (1+\varepsilon) \cdot P_i] \bigcap [C_1 \cdot \hat{\xi}_{i,k-1}, C_2 \cdot \hat{\xi}_{i,k-1}].$$
(4)

This range accounts for both the plausible range and the preceding shape. ε , C_1 , and C_2 are userdefined parameters. (In our experiments, $0 \le \varepsilon \le 0.2$, $0.75 \le C_1 \le 0.9$, and $1.1 \le C_2 \le 1.25$.)

Our goal is to compute the location of each bird in the flock using the above relative power reduction values $\hat{\xi}_{i,k}$. However, for computing these positions, we will show that we must know the absolute power reduction value for each bird, which is unavailable. (It cannot be drawn from the movie, since it is meaningless by itself; only the distribution of the values among the birds provides the shape information.)

To overcome this problem and compute the absolute power reduction values $\hat{e}_{i,k}$, which is defined by

$$\hat{e}_{i,k} = \hat{\xi}_{i,k} \sum_{n=1}^{N} \hat{e}_{n,k},$$
(5)

we take advantage of a special property of the total power reduction $\sum_{n=1}^{N} \hat{e}_{n,k}$ of the flock: According to the displacement theorem of Munk, the total power reduction of flocks having the same number of birds and x-distances is constant [13]. The longitudinal displacements have no influence on it, but rather only on its distribution.

We thus choose arbitrary *y*-positions and assume that the *x*-distance is constant, chosen from the range of possible mean *x*-distances. We now compute the total power reduction value for these positions. Note that though each bird's power reduction depends on this specific *y*-position (and thus cannot be used), the total sum can be utilized in our calculations. Finally, having found the absolute power reduction values $\{\hat{e}_{i,k}\}_{i=1}^{N}$ (Equation 5) corresponding to the relative power reduction values $\{\hat{\xi}_{i,k}\}_{i=1}^{N}$, the positions of the birds are calculated. They are the solution of the following N non-linear equations system:

$$e_i(x_{1i},...,x_{Ni},y_{1i},...,y_{Ni}) = \hat{e}_{i,k} \quad i = 1 \le i \le N,$$
(6)

where $e_i(x_{1i},...,x_{Ni},y_{1i},...,y_{Ni})$ is the function presented in Equation 1, which takes the geometric function g_{ij} into account.

The solution to this system gives us the N-1 unknown relative y-positions, and hence the exact positions of each bird in the flock. The system is solved by the Levenberg-Marquardt algorithm for each key shape.

VI. FLOCK INITIATION

The input to this stage is the *target shape*, which is the first key shape in the steady flight. This shape consists of the positions of the *N* birds. The challenge is to generate a flock initiation that converges to it from any initial conditions.

Energy-savings models cannot be utilized for initiation, since they are applicable only to the stage of steady flight. Instead, we adopt the artificial bird model and specify new rules, whose application for each bird results in convergence of the target shape.

At each time step of the initiation animation, each bird is characterized by its flight direction, flight speed, and position. Its actions, which aim at satisfying its drives, result from the perceived world at the previous time step. Below we discuss all these components.

A. Modeling perception

Artificial animals have senses through which they perceive limited information about the world. Since in nature most line-formation flocks take off from obstacle-free environments and fly in the open sky, we define the world as a collection of birds. We consider the visual perception and apply two operations at every time step: determination of the visible birds and detection of the reference bird (Figure 7).

To determine the visible flockmates, the visual volume is first set (in our setup, it is 90^0 per eye). Then, each bird is modeled by a point and the *Hidden Point Removal (HPR)* operator [15] is applied. This operator computes visibility directly from a point set, by extracting the points



Fig. 7. The perception model of the bird in the blue circle. The shaded area is the visual volume. Only the black birds within this volume are visible, while the white birds are hidden. The reference bird is in a red circle.

that reside on the convex hull of the spherically-inverted point set. This inversion reflects the points with respect to a sphere of radius R. The question is how to choose R so as to take into account the sparsity of the point set and the fact that the points have volume. Our experiments indicated that choosing R to be half of the bird's wingspan provides good results. Recall that the wingspan depends on the specific bird species. For instance, for a Canadian goose it is 1.5 meters.

Next, each bird chooses a *reference bird* from its visible flockmates – a bird it aims to follow. Since following a bird should require as little effort as possible, in terms of speed and direction changes, the reference bird should be nearby. The closest bird within a visual volume of 0.35π per eye is chosen. Only if no such bird exists, the closest visible bird is selected. Thus, at the beginning of the animation several different birds may have the same reference bird, but this will be resolved by the separation drive as the animation proceeds. If a bird does not have any visible flockmates, it does not have a reference bird (although, it may gain one in subsequent time steps). Each bird acquires the following information about its reference bird: speed, direction, relative side (left/right), and whether it is the leader (front-most) of the flock.

B. Modeling drives and actions

Below we describe our proposed drives—*target-reaching* and *separation*—and their realization.

1) *Target-reaching drive:* The target-reaching drive accounts for the main distinction between line-formation flocks and cluster-formation flocks. Rather than attempting to fly "close" to their

14



Fig. 8. Side and distance determination. The birds in the middle preserve the correct distances from their targets, but fail to choose the correct relative side. The birds on the right do not preserve the target distances.

flockmates (by attraction) to form a cluster, the birds try to converge to a certain shape. Since no global decisions are made, this target shape should be attained through the actions of each individual bird. The drive of each bird (excluding the leader) is, hence, to reach a position at a certain *target distance* on a certain *side* (left/right) relative to its reference bird (which can change at every time step).

The resulting actions of this drive are changes in the bird's speed and direction. These actions are performed by first determining the relative side and distance from the reference bird, as follows.

Side determination: Every bird determines its relative side, so as to satisfy global consistency and avoid "zigzagging," as illustrated in Figure 8. At every time step, if the reference bird has an undetermined side (e.g., the first time step), the bird chooses its side according to its current relative position: if it is to the right of the reference bird, its side will be set to the right. Otherwise, the bird's side is set to that of its reference bird. Note that this will result in a shape in which the leader may have followers on both sides, whereas a non-leader may be followed on a single side.

Target-distance determination: A correspondence between each animated bird and a bird of the target shape is determined. First, the leaders are matched. Then, birds whose reference birds have already been matched are also matched consistently. Note that many animated birds may initially be matched to the same bird of the target shape. This will be eventually resolved, as the animation proceeds. The target distances (vector) $\vec{\Delta T_i}$ are derived from the corresponding distances in the target shape.

Calculation of speed and direction: Speed changes depend on three factors: distance to the



Fig. 9. The reference bird is in the red circle; the gray bird is its predicted position at the next time step; the red triangle is the target position, at target-distance ΔT_i from the reference bird's predicted position.

target position, velocity of the reference bird, and the bird's flight direction. Obviously, if the distance is large, the bird should accelerate in order to reduce the gap. Moreover, if the reference bird is moving fast, its followers must adjust their speed to catch up with it. If the flight directions of the reference bird and its follower differ considerably, increased speed might result in increased distance, as a result of a fast flight in a wrong direction. To model the bird's finite energy and avoid inconsistencies, we restrict the maximal changes to both direction and speed.

Let *i* and *j* be the indices of a specific bird and its reference bird, respectively. Let \vec{P}_j be the position of the reference bird, flying at speed v_j in direction $\vec{d}_j = (\sin(\theta_j), \cos(\theta_j))$. We wish to calculate the speed v'_i and direction \vec{d}'_i of bird *i* in the next time frame (Figure 9).

First, the position of the reference bird \vec{P}'_j in the next time step is predicted as $\vec{P}'_j = \vec{P}_j + v_j \vec{d}_j$. Then, the target position of bird *i* is calculated as: $\vec{T}_i = \vec{P}'_j - \vec{U} * \vec{\Delta T}_i$, where $\vec{\Delta T}_i$ is the target distance obtained before and \vec{U} is (1 1) or (-1 1), depending on the bird's side.

The updated flight speed v'_i is calculated as follows. Let r_i be the distance of bird *i* to its target position, $r_i = \|\vec{P}_i - \vec{T}_i\|$, \vec{d}_{iT} be a unit vector in the direction from \vec{P}_i to the target position \vec{T}_i , and Δd be the angle between \vec{d}_i and \vec{d}_{iT} . Then,

$$v'_{i} = v_{j} \cdot \left(1 - \frac{|\Delta d|}{\pi} + K\sqrt{|r_{i} - v_{j}|}\right).$$
(7)

In this equation, the new speed v'_i increases as the reference bird's speed increases. Moreover, it increases with the distance r_i (so as to catch up). However, it decreases if the flight direction is not aimed at the target position. Finally, the sign of parameter K is set according to the relative

The direction angle of the next frame, θ_i' , is derived by:

$$\theta_i' = \theta_i + \alpha \frac{\Delta d}{\max(r_i, 1)}.$$
(8)

Here, the change in direction is inversely proportional to the distance r_i , i.e., when the bird is far from its reference bird, it will change its flight direction slowly, whereas if it is close, it will change its direction quickly, to avoid missing the target. It is also proportional to Δd , the total expected change in angle. α is a constant that is set to 0.2 in our experiments.

2) Separation drive: The separation drive has two goals. First, it ensures that at the end of the initiation, each bird (except the leader) is followed by at most one bird. In addition, it aids in avoiding collisions between the birds.

This drive is realized as follows. Each bird detects the birds having the same reference bird among its visible flockmates. If such a bird is found and the distance between them is sufficiently small, it will choose a new reference bird. Hence, if two birds have the same reference bird, only the rear bird will change its reference bird. If the distance between the birds is large, they will continue in their flight until they come closer to each other. (The threshold used is ten times the wingspan.)

Note that our separation drive is fundamentally different from those of [4], [24]. While the latter simulates repulsion from visible flockmates, ours simulates repulsion from the "cluster" of flockmates following the same reference bird. In addition, rather than resulting in changes in speed and direction, our drive results is selecting a new reference bird.

VII. RESULTS

The accompanying movie and Figures 10–14 show some results. Figure 10 shows an example where the number of birds in the example movie and the animation is the same. The animated birds are super-imposed on the real barnacle geese, so that they could be compared. While the shapes are not identical, they bear similarity and look as natural as that of the real flock.

Figures 11–12 show snapshots from an animation that uses the same example movie (of 9 birds), yet creates an animation of 14 birds. Figure 12 demonstrates initiation from random



Fig. 10. Snapshots from a steady flight of 9 barnacle geese. The brown (higher) birds are the animated ones, whereas the gray ones belong to the original movie (along with the original background). Snapshots from the initiation of this flock are presented in Figure 1, using a different background.



Fig. 11. Snapshots from a steady flight animation of a 14-bird flock, whose behavior was learned from the movie of the 9-bird flock of Figure 10.



Fig. 12. Snapshots from the initiation of a 14-bird flock, whose steady flight is shown in Figure 11.

locations, whereas Figure 11 shows the steady flight stage. It can be seen that the general shapes are maintained, creating realistic-looking flocks. For instance, the two rear birds on the right side are very close (in x-coordinates) to each other. This happens in the original movie for the rear birds on the left. Moreover, it is interesting to note that the flock changes from a U-shape to a V-shape. This occurs both in the original 9-bird flock and in the animated 14-bird flock.

Figure 13 demonstrates snapshots from an animation of snow geese. It compares the original movie (bottom) to our animation (top). Again, our data-driven approach is capable of reconstruct-



Fig. 13. Snapshots from a steady flight animation of a flock of 16 snow geese. These frames show both the original movie (bottom birds) and the animated birds (on top).



Fig. 14. Snapshots from an animation of 41 cranes (top) compared to the real movie (bottom). Note the interesting phenomenon of a bird switching sides.

ing similar general shapes. Figure 14 demonstrates an animation of a large flock of 41 cranes. Our animation manages to construct the V-shape of the flock. In the animation, in contrast to the example, the right side of the flock is longer than the left, and therefore the bird switches position in the opposite direction. In nature, birds switching sides during the flight is pretty rare. Our animation-by-example approach manages to animate it.

Data acquisition: Data acquisition is an interesting problem in its own right, but outside the scope of the current paper. In our examples, we tried to choose example movies in which the flocks were recorded (almost) from above. The camera motion and the depth of the birds were not estimated, similarly to [18]. The frames are manually chosen from the input movie, and the birds are marked. This takes about 2-5 minutes of manual work per movie.

VIII. CONCLUSION

This paper has introduced an approach for animating flocks of birds flying in line-formation patterns. We distinguish between the behavior of such flocks during initiation and their behavior during the steady flight and propose different techniques for modeling and animating each of these stages. A data-driven approach that utilizes an energy-savings model is proposed for animating the steady flight. We show that this problem can be formulated as a system of non-linear equations and solved. Rules are used to govern the behavior of the birds during initiation. These rules differ from those of previous work, which have focused on flocks flying in clusters.

Our results are evaluated in two manners. First, they are compared to real flocks, which demonstrate that they look natural for a variety of line-formation patterns. Second, we guarantee that the energy-saving values are reasonable for each bird in the flock. Since it is believed that these values are the major cause for flying in formation, this assures that our flocks are indeed plausible.

Our method has several limitations, which should be addressed in the future. First, our initiation algorithm cannot handle inverted shapes (such as inverted U or V) that exist in nature. These shapes have two leaders, while our technique assumes a single leader. Similarly, some flocks are shaped as a large V with another line "attached" to the middle of one of its sides. Our techniques cannot simulate such flocks. Second, we do not specifically model environmental conditions, such as wind gusts. However, since our animation is guided by a real-world example, the influence of some environmental conditions is indirectly accounted for. Third, in the data acquisition stage we assume that the example movies are shot from above/below. Instead, the method should consider perspective transformation. Finally, more advanced techniques for animating wing flapping, such as [33], can be used.

REFERENCES

- M. Anderson, E. McDaniel, and S. Chenney. Constrained animation of flocks. In ACM symposium on Computer animation, pages 286–297, 2003.
- M. Andersson and J. Wallander. Kin selection and reciprocity in flight formation? *Behavioral Ecology*, 15(1):158–162, 2004.
- [3] J.P. Badgerow and F.R. Hainsworth. Energy savings through formation flight? A reexamination of the vee formation. J. of Theoretical Biology, 93(1):41–52, 1981.
- [4] I.L. Bajec, N. Zimic, and M. Mraz. Simulating flocks on the wing: the fuzzy approach. *Journal of Theoretical Biology*, 233(2):199–220, 2005.

- [5] C. Cutts. Energy savings in formation flight of pink-footed geese. J. of Experimental Biology, 189(1):251-261, 1994.
- [6] M.A. Fischler and R.C. Bolles. Random sample consensus: a paradigm for model fitting with applications to image analysis and automated cartography. *Comm. of the ACM*, 24(6):381–395, 1981.
- [7] G.W. Flake. The computational beauty of nature: Computer explorations of fractals, chaos, complex systems, and adaptation. MIT Pr., 1998.
- [8] L.L. Gould and F.H. Heppner. The vee formation of Canada geese. Auk, 91:494–506, 1974.
- [9] F.R. Hainsworth. Induced drag savings from ground effect and formation flight in brown pelicans. J. of Experimental Biology, 135(1):431-444, 1998.
- [10] F.H. Heppner. Avian flight formations. Bird-Banding, 45(2):160-169, 1974.
- [11] F.H. Heppner. Three-dimensional structure and dynamics of bird flocks. *Animal Groups in Three Dimensions*, pages 68–89, 1997.
- [12] F.H. Heppner and U. Grenander. A stochastic nonlinear model for coordinated bird flocks. *American Assoc. for the Advancement of Science*, 1990.
- [13] D. Hummel. Aerodynamic aspects of formation flight in birds. Journal of Theoretical Biology, 104(3):321-347, 1983.
- [14] D. Hummel. Formation flight as an energy-saving mechanism. Israel J. Zoology, 41(3):261–278, 1995.
- [15] S. Katz, A. Tal, and R. Basri. Direct visibility of point sets. ACM Trans. on Graph., 26(3):24:1-11, 2007.
- [16] R.M. King and A. Gopalarathnam. Ideal aerodynamics of ground effect and formation flight. *Journal of Aircraft*, 42(5):1188, 2005.
- [17] K.H. Lee, M.G. Choi, Q. Hong, and J. Lee. Group behavior from video: a data-driven approach to crowd simulation. In Symp. on Comp. Animation, pages 109–118, 2007.
- [18] A. Lerner, Y. Chrysanthou, and D. Lischinski. Crowds by example. Graphics Forum, 26(3), 2007.
- [19] P.B.S Lissaman and C.A. Shollenberger. Formation flight of birds. Science, 168(3934):1003–1005, 1970.
- [20] R.A. Metoyer and J.K. Hodgins. Reactive pedestrian path following from examples. *The Visual Computer*, 20(10):635–649, 2004.
- [21] S.R. Musse and D. Thalmann. A model of human crowd behavior: Group inter-relationship and collision detection analysis. In *Computer Animation and Simulation*, pages 39–51, 1997.
- [22] A. Nathan and V.C. Barbosa. V-like formations in flocks of artificial birds. Artificial Life, 14(2):179–188, 2008.
- [23] A. Okubo. Dynamical aspects of animal grouping: swarms, schools, flocks, and herds. *Advances in Biophysics*, 22:1, 1986.
- [24] C.W. Reynolds. Flocks, herds and schools: A distributed behavioral model. SIGGRAPH Comput. Graph., 21(4):25–34, 1987.
- [25] C.W. Reynolds. Steering behaviors for autonomous characters. In Game Developers Conference, pages 763–782, 1999.
- [26] C.M. Sewatkar, A. Sharma, and A. Agrawal. A first attempt to numerically compute forces on birds in v formation. *Artificial Life*, 16(3):245–258, 2010.
- [27] W. Shao and D. Terzopoulos. Autonomous pedestrians. Graphical Models, 69(5-6):246-274, 2007.
- [28] T. Sugimoto. A theoretical analysis of formation flight as a nonlinear self-organizing phenomenon. IMA J. of Applied Mathematics, 68(5):441–470, 2003.
- [29] M. Sung, M. Gleicher, and S. Chenney. Scalable behaviors for crowd simulation. In *Computer Graphics Forum*, volume 23, pages 519–528, 2004.
- [30] A. Treuille, S. Cooper, and Z. Popović. Continuum crowds. ACM Transactions on Graphics, 25(3):1160–1168, 2006.

- [31] X. Tu and D. Terzopoulos. Artificial fishes: physics, locomotion, perception, behavior. In SIGGRAPH, pages 43-50, 1994.
- [32] H. Weimerskirch, J. Martin, Y. Clerquin, P. Alexandre, and S. Jiraskova. Energy saving in flight formation. *Nature*, 413(6857):697–8, 2001.
- [33] J. Wu and Z. Popović. Realistic modeling of bird flight animations. In *ACM Transactions on Graphics (TOG)*, volume 22, pages 888–895. ACM, 2003.